CVS specialty

Rheumatology Enrollment Form Medications A (Actemra®)

Fax Referral To: 1-877-232-5455 Address: 500 Ala Moana Blvd., Ste 1-A Honolulu, HI 96813 Phone: 1-800-896-1464

	INFORMATION (Con	Six Simple Steps to		lerral	
				City, State, ZIP:	
Preferred Conta	act Methods: Phone (to r	/(durced	Text (to cell # p	rovided below) Email (to end	mail provided below)
				acy will attempt to contact by	
	• • • • •		· · ·	Gender: 🗌 Mal	-
	/	Last Four of SSI	<u> </u>	Primary Language:	
	BER INFORMATION				
			ate License #:		
NPI #:	DEA #:	Group or Hosp	ital:		
Address:		Č	ity, State, ZIP:		
Phone:	Fax	Contact	Person:	Contact's Phone:	
				e cards with this form, if avail	
	SIS AND CLINICAL	NFORMATION			
	Ship to: Patie				
Diagnosis (ICE					
	matoid Arthritis, Unspecifie	d	☐ M45.9 Ank	ylosing Spondylitis of Unspec	ified Sites in Spine
	opathic Psoriasis, Unspecif			er Psoriatic Arthropathy	
				e: Description	
	CD-10 information, please v	•			
	specialty.com/wps/portal/sp				
Patient Clinica					
		Weight: Ib/kg	Height: in/cr	n TB Test Result:	Date:
Nursing:		00	J		
Specialty pharn	nacy to coordinate injection	training/ home health infu	sion nurse visit nec	essary 🗌 Yes 🔲 No	
Site of Care:	MD office 🔲 Infusion Clin	ic 🔲 Outpatient Health [Home Health	· — —	
Injection training	g not necessary. Date traini	ng occurred:			
Reason: M	D office training patient 🗌 F	Pt already independent	Referred by MD to	alternate trainer	
5 PRESCRI	PTION INFORMATI	ON			
MEDICATION	STRENGTH	D	OSE & DIRECTION	NS Q	UANTITY/REFILLS
	🗌 80 mg/4 mL	Induction Dose: Infu			Quantity:
Actemra	200 mg/10 mL	Maintenance Dose:	Infuse 8 mg/kg eve	ry 4 weeks.	Refills:
	🗌 400 mg/20 mL	Other:			
	162mg/0.9 mL prefilled			62mg SC every other week,	Quantity:
Actemra	syringe	followed by an increase			Refills:
Patient is interested	in patient support programs	For patients weighin	$\underline{IG} \ge 100 \underline{KG}$: Inject 1 E NOT ALLOWED	62mg SC every week. Ancillary supplies and kits provide	ded as needed for administratio
		HYSICIAN SIGN			
		(Date)	DISPENSE AS WRITT		(Date)
Y		()			(Date)
^		·····	^		
The information pr	ovided above is true and accurate	a to the best of my knowledge	with supporting docum	entation in the nationt's medical rec	ord By signing below I

uye hereby authorize CVS Specialty Pharmacy and/or its affiliate pharmacies to complete and submit prior authorization (PA) requests to payors for the prescribed medication for this patient and to attach this Enrollment Form to the PA request as my signature.

CONFIDENTIALITY NOTICE: This communication and any attachments may contain confidential and/or privileged information for the use of the designated recipients named above. If you are not the intended recipient, you are hereby notified that you have received this communication in error and that any review, disclosure, dissemination, distribution or copying of it or its contents is prohibited. If you have received this communication in error, please notify the sender immediately by telephone and destroy all copies of this communication and any attachments.

Plan member privacy is important to us. Our employees are trained regarding the appropriate way to handle members' private health information.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Pharmacy, Inc. or one of its affiliates.

Medications A-C Rheumatology Enrollment Form

(Avsola[™]Cimzia®,Cosentyx®)

Please complete Patient and Prescriber information						
Patient Name: Patient DOB:						
Prescriber Nam	e:	Prescriber Phone:	_			
5 PRESCR	5 PRESCRIPTION INFORMATION					
MEDICATION	STRENGTH	DOSE & DIRECTIONS	QUANTITY/REFILLS			
Avsola	100 mg vial	☐ Rheumatoid Arthritis <u>Induction Dose: In conjunction with</u> <u>methotrexate</u> Infuse IV at 3 mg/kg (Dose =mg) at weeks 0, week 2, week 6 and every 8 weeks thereafter. ☐ Rheumatoid Arthritis <u>Maintenance Dose:</u> Infuse 3 mg/kg every 8 weeks.	Quantity: # of 100 mg vial Refills:			
		 Psoriatic Arthritis Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, week 2, week 6 and every 8 weeks thereafter. Psoriatic Arthritis Maintenance Dose: Infuse 5 mg/kg every 8 weeks. Ankylosing Spondylitis Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, week 2, week 6 and every 6 weeks thereafter. Ankylosing Spondylitis Maintenance Dose: Infuse 5 mg/kg every 6 weeks. Other: 				
🗌 Cimzia	Cimzia Starter Kit (6 prefilled syringes)	Induction Dose: 400 mg initially and at week 2 and 4, (given as 2 SC of 200 mg each) followed by 200 mg every other week;	Quantity: 1 Kit Refills: 0			
🗌 Cimzia	☐ 200mg/1 mL prefilled syringe ☐ 200mg vial	 <u>Maintenance Dose</u>: Inject 200mg SC every OTHER week. <u>Maintenance Dose</u>: Inject 400mg SC every four weeks. ☐ Other: 	Quantity: Refills:			
Cosentyx	 ☐ Sensoready® pen 150 mg/mL injection ☐ Prefilled syringe 150 mg/mL injection 	Psoriatic Arthritis with Coexistent Moderate to Severe Plaque Psoriasis Loading Dose: Inject 300 mg (two injections) SC at weeks 0, 1, 2, 3 and 4. Maintenance Dose: Inject 300 mg (two injections) SC every 4 weeks. Other Psoriatic Arthritis or Ankylosing Spondylitis With Loading Dose: Inject 150 mg (one injection) SC at weeks 0, 1, 2, 3 and 4, and then every 4 weeks thereafter. Without Loading Dose: Inject 150 mg (one injection) SC every 4 weeks. Other:	Quantity: Refills:			
Patient is interested in patient support programs STAMP SIGNATURE NOT ALLOWED Ancillary supplies and kits provided as needed for administration BPHYSICIAN SIGNATURE REQUIRED						

PRODUCT SUBSTITUTION PERMITTED

(Date)

DISPENSE AS WRITTEN X

(Date)

Х

The information provided above is true and accurate to the best of my knowledge, with supporting documentation in the patient's medical record. By signing below, I hereby authorize CVS Specialty Pharmacy and/or its affiliate pharmacies to complete and submit prior authorization (PA) requests to payors for the prescribed medication for this patient and to attach this Enrollment Form to the PA request as my signature.

CONFIDENTIALITY NOTICE: This communication and any attachments may contain confidential and/or privileged information for the use of the designated recipients named above. If you are not the intended recipient, you are hereby notified that you have received this communication in error and that any review, disclosure, dissemination, distribution or copying of it or its contents is prohibited. If you have received this communication in error, please notify the sender immediately by telephone and destroy all copies of this communication and any attachments.

Plan member privacy is important to us. Our employees are trained regarding the appropriate way to handle members' private health information.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Pharmacy, Inc. or one of its affiliates.

Medications E-I Rheumatology Enrollment Form

(Enbrel®, Humira®, Ilaris®, Inflectra®)

Please complete Patient and Prescriber information Patient Name: Patient DOB: Prescriber Phone: Prescriber Name: 5 PRESCRIPTION INFORMATION MEDICATION STRENGTH QUANTITY/REFILLS **DOSE & DIRECTIONS** Quantity: 25mg/0.5 mL prefilled syringe Refills: 25mg vial 50mg/mL Sureclick™ Autoinjector 50mg/mL prefilled ☐ Inject 25mg SC TWICE a week (72 – 96 hours apart). syringe Enbrel 50 mg/mL Enbrel Inject 50mg SC ONCE a week. Mini[™] prefilled cartridge Other: for use with the AutoTouch[™] reusable autoinjector only (Prescriber MUST supply). CVS does not order the autoinjector. Quantity: ☐ 40 mg/0.4 mL Pen ☐ Inject 40mg SC every OTHER week. Citrate Free Refills: Humira Other: 40 mg/0.4 mL Prefilled Syringe Citrate Free For patients weighing ≥ 7.5 kg: Inject 4 mg/kg (with a maximum of Quantity: 300 mg) SC every 4 weeks. Each single-dose vial of ILARIS Refills: 150 mg/mL injection Ilaris (canakinumab) Injection delivers 150 mg/mL sterile, preservativesolution free, clear to slightly opalescent, colorless to a slight brownish to yellow solution. Rheumatoid Arthritis Induction Dose: In conjunction with Quantity: methotrexate Infuse IV at 3 mg/kg (Dose = mg) at weeks 0, # of 100 mg vial week 2, week 6 and every 8 weeks thereafter. Refills: Rheumatoid Arthritis Maintenance Dose: Infuse 3 mg/kg every 8 weeks. Psoriatic Arthritis Induction Dose: Infuse IV at 5 mg/kg (Dose = mg) at weeks 0, week 2, week 6 and every 8 weeks thereafter. ☐ Inflectra 100 mg vial Psoriatic Arthritis Maintenance Dose: Infuse 5 mg/kg every 8 weeks. Ankylosing Spondylitis Induction Dose: Infuse IV at 5 mg/kg (Dose = mg) at weeks 0, week 2, week 6 and every 6 weeks thereafter. Ankylosing Spondylitis Maintenance Dose: Infuse 5 mg/kg every 6 weeks. Other: STAMP SIGNATURE NOT ALLOWED Ancillary supplies and kits provided as needed for administration Patient is interested in patient support programs **6 PHYSICIAN SIGNATURE REQUIRED**

PRODUCT SUBSTITUTION PERMITTED	(Date)	DISPENSE AS WRITTEN	(Date)
X		Χ	

The information provided above is true and accurate to the best of my knowledge, with supporting documentation in the patient's medical record. By signing below, I hereby authorize CVS Specialty Pharmacy and/or its affiliate pharmacies to complete and submit prior authorization (PA) requests to payors for the prescribed medication for this patient and to attach this Enrollment Form to the PA request as my signature.

CONFIDENTIALITY NOTICE: This communication and any attachments may contain confidential and/or privileged information for the use of the designated recipients named above. If you are not the intended recipient, you are hereby notified that you have received this communication in error and that any review, disclosure, dissemination, distribution or copying of it or its contents is prohibited. If you have received this communication in error, please notify the sender immediately by telephone and destroy all copies of this communication and any attachments.

Plan member privacy is important to us. Our employees are trained regarding the appropriate way to handle members' private health information.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Pharmacy, Inc. or one of its affiliates.

Medications K-O Rheumatology Enrollment Form

(Kevzara®,Olumiant®,Orencia®, Otezla®)

	Please	e complete Patient and Prescriber information	
Patient Name:		Patient DOB:	_
Prescriber Name: Prescriber Phone:			_
5 PRESCRI	PTION INFORMATIO	N	
MEDICATION	QUANTITY/REFILLS		
🗌 Kevzara	☐ 200 mg/1.14 mL prefilled syringe (pk of 2) ☐ 150 mg/1.14 mL prefilled syringe (pk of 2) ☐ 200 mg/1.14 mL prefilled pen (pk of 2) ☐ 150 mg/1.14 mL prefilled pen (pk of 2)	 Inject 200 mg SC once every two weeks. Inject 150 mg SC once every two weeks. 	Quantity: Refills:
Olumiant	2 mg tablet 1 mg tablet	☐ Take 2 mg PO once daily ☐ Other:	Quantity: Refills:
🗌 Orencia	☐ 125mg prefilled syringe ☐ ClickJect Autoinjector 125 mg/mL pack of 4	 Inject 125mg SC every week <u>After Single IV Loading Dose</u>: Inject 125mg SC within a day and 125mg SC every week thereafter. <u>Patients Unable to Receive an IV Loading Dose</u>: Inject 125 mg SC every week. <u>Patients Transitioning from IV Infusion Therapy</u>: Inject 125 mg SC instead of the next scheduled IV dose, followed by 125mg SC injections every week thereafter. 	Quantity: Refills:
Orencia	250 mg vial	☐ Infuse mg at weeks 0, 2 and 4, then every 4 weeks thereafter. ☐ Other:	Quantity: Refills:
🗌 Otezla	Titration Starter Pack	Day 1: 10 mg PO in the morning. Day 2: 10 mg PO in the morning and 10 mg PO in the evening. Day 3: 10 mg PO in the morning and 20 mg PO in the evening. Day 4: 20 mg PO in the morning and 20 mg PO in the evening. Day 5: 20 mg PO in the morning and 30 mg PO in the evening. Day 6 and thereafter: 30 mg PO twice daily.	Quantity: 1 Pack Refills: 0
🗌 Otezla	30 mg tablet	<u>Maintenance Dose:</u> 30 mg PO twice daily. Other:	Quantity: Refills:

Patient is interested in patient support programs

6 PHYSICIAN SIGNATURE NOT ALLOWED Ancillar

Ancillary supplies and kits provided as needed for administration

PRODUCT SUBSTITUTION PERMITTED	(Date)	DISPENSE AS WRITTEN	(Date)
X		X	

The information provided above is true and accurate to the best of my knowledge, with supporting documentation in the patient's medical record. By signing below, I hereby authorize CVS Specialty Pharmacy and/or its affiliate pharmacies to complete and submit prior authorization (PA) requests to payors for the prescribed medication for this patient and to attach this Enrollment Form to the PA request as my signature.

CONFIDENTIALITY NOTICE: This communication and any attachments may contain confidential and/or privileged information for the use of the designated recipients named above. If you are not the intended recipient, you are hereby notified that you have received this communication in error and that any review, disclosure, dissemination, distribution or copying of it or its contents is prohibited. If you have received this communication in error, please notify the sender immediately by telephone and destroy all copies of this communication and any attachments.

Plan member privacy is important to us. Our employees are trained regarding the appropriate way to handle members' private health information.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Pharmacy, Inc. or one of its affiliates.

Medications R-S Rheumatology Enrollment Form (Remicade®, Renflexis®, Rinvoq®, Rituxan®, Simponi®)

Detient Name:		e complete Patient and Prescriber information	
Patient Name: Prescriber Name		Patient DOB: Prescriber Phone:	-
			-
MEDICATION	STRENGTH	DOSE & DIRECTIONS	QUANTITY/REFILLS
		■ Rheumatoid Arthints <u>Induction Dose</u> . In conjunction with <u>methotrexate</u> Infuse IV at 3 mg/kg (Dose =mg) at weeks 0, week 2, week 6 and every 8 weeks thereafter. ☐ Rheumatoid Arthritis <u>Maintenance Dose</u> : Infuse 3 mg/kg every 8 weeks.	Quantity: Refills:
☐ Remicade	100 mg vial	 Psoriatic Arthritis Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, week 2, week 6 and every 8 weeks thereafter. Psoriatic Arthritis Maintenance Dose: Infuse 5 mg/kg every 8 weeks. Ankylosing Spondylitis Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, week 2, week 6 and every 6 weeks thereafter. Ankylosing Spondylitis Maintenance Dose: Infuse 5 mg/kg every 6 weeks. 	
Renflexis	100 mg vial	 Other:	Quantity: # of 100 mg vial Refills:
		thereafter. Ankylosing Spondylitis <u>Maintenance Dose: Infuse 5 mg/kg</u> every 6 weeks. Other:	
🗌 Rinvoq	15 mg	Take one 15 mg tablet PO once daily. Other:	Quantity: Refills:
☐ Rituxan	☐ 100 mg/10 mL vial ☐ 500 mg/50 mL vial	 Infuse two doses of 1000 mg separated by 2 weeks. Other: 	Quantity: Refills:
Simponi	50mg/0.5mL prefilled SmartJect® Autoinjector 50mg/0.5mL prefilled syringe	Inject 50mg SC once a month. Other:	Quantity: Refills:
Patient is interested in	patient support programs		ded as needed for administration
PRODUCT SUBSTITU X		(Date) DISPENSE AS WRITTEN X	(Date)

The information provided above is true and accurate to the best of my knowledge, with supporting documentation in the patient's medical record. By signing below, I hereby authorize CVS Specialty Pharmacy and/or its affiliate pharmacies to complete and submit prior authorization (PA) requests to payors for the prescribed medication for this patient and to attach this Enrollment Form to the PA request as my signature.

CONFIDENTIALITY NOTICE: This communication and any attachments may contain confidential and/or privileged information for the use of the designated recipients named above. If you are not the intended recipient, you are hereby notified that you have received this communication in error and that any review, disclosure, dissemination, distribution or copying of it or its contents is prohibited. If you have received this communication in error, please notify the sender immediately by telephone and destroy all copies of this communication and any attachments.

Plan member privacy is important to us. Our employees are trained regarding the appropriate way to handle members' private health information.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Pharmacy, Inc. or one of its affiliates.

Medications S-Z Rheumatology Enrollment Form (Simponi ARIA®, Stelara®, Taltz®,Tremfya®,Xeljanz®)

Please complete Patient and Prescriber information

Patie	ent D	OB:
_		

Patient Name: Prescriber Name:

Prescriber Phone:

5 PRESCRIPTION INFORMATION					
MEDICATION	S	TRENGTH		DOSE & DIRECTIONS	QUANTITY/REFILLS
☐ Simponi ARIA	50 mg/4 mL in a single use vial			Infuse 2 mg/kg over 30 minutes at weeks 0 and 4, then every weeks thereafter.	/ 8 Quantity: # of 50 mg vial Refills:
Stelara	☐ 45mg/0.5mL prefilled syringe ☐ 90mg/mL prefilled syringe			 □ For patients weighing ≤100 kg (220 lbs): Inject 45 mg SC initially and 4 weeks later, followed by 45 mg every 12 weeks □ For patients weighing >100 kg (220 lbs): Inject 90 mg SC initially and 4 weeks later, followed by 90 mg every 12 weeks □ Other:	
☐ Taltz	 80 mg Single Dose Autoinjector 80 mg Single Dose Prefilled Syringe 			Psoriatic Arthritis with Coexistent Moderate to Severe Plaque Psoriasis Dosing: Starting Dose: Inject SC two 80 mg injections on Day 1, the begin the induction dose 2 weeks later. Induction Dose: Inject SC one 80 mg injection every 2 week (weeks 2, 4, 6, 8, 10, and 12). Maintenance Dose: Inject SC one 80 mg injection every 4 weeks.	nen 3 Pens/Syringes 2 Pens/Syringes eks 1 Pens/Syringes Refills:
🗌 Taltz	 80 mg Single Dose Autoinjector 80 mg Single Dose Prefilled Syringe 			Psoriatic Arthritis Dosing and Ankylosing Spondylitis Dosing: Starting Dose: Inject SC two 80 mg injections on Day 1. Maintenance Dose: Inject SC one 80 mg injection every 4 weeks. Non-radiographic Axial Spondyloarthritis Dosing: Dose: Inject SC one 80 mg injection every 4 weeks	Quantity: 2 Pens/Syringes 1 Pens/Syringes Refills:
Tremfya	100 mg/mL prefilled syringe		d	Psoriatic Arthritis Dosing: 100mg administered by SC at Week 0, Week 4 and every weeks thereafter	8 Quantity: Refills:
🗌 Xeljanz	5 mg Tablet			Take one 5 mg tablet PO twice daily Take one 11 mg PO once daily Other:	Quantity: Refills:
			Home I	nfusion/Coram AIS: DOSE/STRENGTH/DIRECTIONS	
		mainta PIV – PORT	ter Care/Flush – Only on drug admin days – SASH or PRN to in IV access and patency NS 5ml (Heparin 10 units/ml 3-5ml if multiple days) /PICC – NS 10ml & Heparin 100units/ml 3-5ml, and/or 10ml sto to access port a cath	QUANTITY/REFILLS Quantity: Refills:	
Epinephrine			Adu Peo Infa PRN s	ult 1:1000, 0.3mL (>30kg/>66lbs) ds 1:2000, 0.3mL (15-30kg/33-66lbs) ant 0.1mL/0.1mL, 0.1mL (7.5-15kg/16.5-33lbs) evere allergic reaction – Call 911 speat in 5-15 minutes as needed	Quantity: Refills:
Patient is interested in			6 PH	STAMP SIGNATURE NOT ALLOWED Ancillary supplies a YSICIAN SIGNATURE REQUIRED	and kits provided as needed for administration
PRODUCT SUBSTITUTION PERMITTED			(Date) DISPENSE AS WRITTEN	(Date)	

Х

The information provided above is true and accurate to the best of my knowledge, with supporting documentation in the patient's medical record. By signing below, I hereby authorize CVS Specialty Pharmacy and/or its affiliate pharmacies to complete and submit prior authorization (PA) requests to payors for the prescribed medication for this patient and to attach this Enrollment Form to the PA request as my signature. CONFIDENTIALITY NOTICE: This communication and any attachments may contain confidential and/or privileged information for the use of the designated recipients

Х

named above. If you are not the intended recipient, you are hereby notified that you have received this communication in error and that any review, disclosure, dissemination, distribution or copying of it or its contents is prohibited. If you have received this communication in error, please notify the sender immediately by telephone and destroy all copies of this communication and any attachments.

Plan member privacy is important to us. Our employees are trained regarding the appropriate way to handle members' private health information.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Pharmacy, Inc. or one of its affiliates.