

Fax Referral To: 1-855-297-1270

Phone: 1-888-280-1190 Address: 280 Avenida Jesus T. Pinero Ste B Rio Piedras, PR 00927 NCPDP: 4026325

	ODMATION (C.)	ix Simple Steps to Subm			
Patient Name:		ete or include demographic she			
				Gender: 🗌 Ma	ıle 🔲 Female
ddress:			State, ZIP Code:		······································
		imary # provided below) 🔲 Text he phone number(s) and email addi			
		ne phone number(s) and email addi ut your prescription(s), account, and			
		tharmacy will attempt to contact by		ru uata rates appty. Mes	sage frequency varies.
mail:		Last Four of	SSN: Pi	rimary Language:	
		st, First):			
	RINFORMATION		Total and the part		
		:	Stata Licanca #:		
DI#	;	Group or Hospital:			
rı #	_ DLA # G	City State 71			
none:		City, State, ZII Contact Person:	Code	ontact's Phone:	
		ase fax copy of prescription and	a insurance cards	with this form, if availa	able (front and back)
	AND CLINICAL INF				
eeds by Date:		Ship to: 🗌 Patient 🗌 Of	fice 🔛 Other:		
iagnosis (ICD-	<u>10):</u>				
L28.1 Prurigo No L40.4 Guttate Ps	odularis 🔲 L40.	0 Psoriasis Vulgaris	L40.1 Generalized P	ustular Psoriasis	
		50 Arthropathic Psoriasis, Unspe			
] L40.54 Juvenile	psoriatic arthritis L40.	59 Other Psoriatic Arthropathy	L40.8 Othe	er Psoriasis	
L40.9 Psoriasis,	Unspecified				
L73.2 Hidradenit	tis Suppurativa	Other Code: Desc	ription:		
atient Clinical	<u>Information:</u>				
lergies:					
/eight:	lb/kg Height:	In/cm TB	Test Result:	Date:	
rior therapy, treatr	ment dates, and reason(s)) for discontinuation:			
reatment status: 🛚	🗌 New to therapy 🔲 Cor	ntinuation of therapy; date of last	treatment//	Needs by date: _	
lursing and Adı	<u>ministration:</u>				
pecialty pharmacy	to coordinate home heal	lth Infusion nurse visit as necessa	ary? 🗌 Yes 🔲 No		
ite of Care: 🔲 Ho			. 🗆	Off: **	
	ome Infusion* 🔲 Coram	n Ambulatory Infusion Suite (AIS)	* 🔛 Prescriber's (Office^^ Utner Infl	usion Clinic
or Remicade/Ren		n Ambulatory Infusion Suite (AIS) three doses to be given in conti		Office^^ Uther Infl	usion Clinic
	nicade Biosimilars: First		olled setting.		usion Clinic
Home Infusion/Co	nicade Biosimilars: First ram AIS: Diluents, Flushe	three doses to be given in conti	rolled setting. drug administration		usion Clinic
Home Infusion/Co Prescriber's Office	nicade Biosimilars: First ram AIS: Diluents, Flushee/Other Infusion Clinic: D	three doses to be given in controls, Supplies, Nursing Services for orug only for facility administratio	rolled setting. drug administration		usion Clinic
Home Infusion/Co Prescriber's Office PRESCRIPTION	nicade Biosimilars: First ram AIS: Diluents, Flushe e/Other Infusion Clinic: D ON INFORMATION	three doses to be given in controls, Supplies, Nursing Services for brug only for facility administratio	r olled setting. drug administration n		
Home Infusion/Co Prescriber's Office PRESCRIPTION	nicade Biosimilars: First ram AIS: Diluents, Flushee/Other Infusion Clinic: D	three doses to be given in controls, Supplies, Nursing Services for brug only for facility administration	rolled setting. drug administration		QUANTITY/REFIL
Home Infusion/Co Prescriber's Office PRESCRIPTION MEDICATION	nicade Biosimilars: First ram AIS: Diluents, Flushe e/Other Infusion Clinic: DON INFORMATION STRENGTH	three doses to be given in controls, Supplies, Nursing Services for brug only for facility administratio DOSE Inject 40 mg SC every week	rolled setting. drug administration n & DIRECTIONS		QUANTITY/REFIL
Home Infusion/Coordinates Office PRESCRIPTION Adalimumab	nicade Biosimilars: First ram AIS: Diluents, Flushe e/Other Infusion Clinic: D ON INFORMATION	three doses to be given in contres, Supplies, Nursing Services for brug only for facility administration DOSE Inject 40 mg SC every week Inject 40 mg SC every other	rolled setting. drug administration n & DIRECTIONS week		QUANTITY/REFIL ☐ 28 days
Home Infusion/Conferescriber's Office PRESCRIPTION MEDICATION Adalimumabaaacf (Unbranded	nicade Biosimilars: First ram AIS: Diluents, Flushe e/Other Infusion Clinic: DON INFORMATION STRENGTH	three doses to be given in controls, Supplies, Nursing Services for brug only for facility administratio DOSE Inject 40 mg SC every week	rolled setting. drug administration n & DIRECTIONS week week	n/therapy teach train.	QUANTITY/REFIL
Home Infusion/Contract Prescriber's Office PRESCRIPTION Adalimumabaacf (Unbranded	nicade Biosimilars: First ram AIS: Diluents, Flushe e/Other Infusion Clinic: DON INFORMATION STRENGTH	three doses to be given in contres, Supplies, Nursing Services for brug only for facility administratio DOSE Inject 40 mg SC every week Inject 40 mg SC every other Inject 80 mg SC every other	rolled setting. drug administration n & DIRECTIONS week week	n/therapy teach train.	QUANTITY/REFIL 28 days 84 days
Home Infusion/Coopersoriber's Office PRESCRIPTION MEDICATION Adalimumabaaacf (Unbranded	nicade Biosimilars: First ram AIS: Diluents, Flushe e/Other Infusion Clinic: DON INFORMATION STRENGTH	three doses to be given in contres, Supplies, Nursing Services for brug only for facility administration. DOSE Inject 40 mg SC every week Inject 40 mg SC every other Inject 80 mg SC every other Inject 80 mg Day 1, followed week after initial dose Inject 40 mg SC every week	rolled setting. drug administration E & DIRECTIONS week week by 40 mg every other	n/therapy teach train.	QUANTITY/REFIL 28 days 84 days
Home Infusion/Conference I	nicade Biosimilars: First ram AIS: Diluents, Flushe e/Other Infusion Clinic: DON INFORMATION STRENGTH	three doses to be given in contres, Supplies, Nursing Services for brug only for facility administration. DOSE Inject 40 mg SC every week Inject 40 mg SC every other Inject 80 mg SC every other Inject 80 mg Day 1, followed week after initial dose	rolled setting. drug administration E & DIRECTIONS week week by 40 mg every other	n/therapy teach train.	QUANTITY/REFIL 28 days 84 days
Home Infusion/Contraction Prescriber's Office PRESCRIPTION Adalimumabaacf (Unbranded Idacio) Adalimumabach	nicade Biosimilars: First ram AIS: Diluents, Flushe e/Other Infusion Clinic: D ON INFORMATION STRENGTH 40 mg/0.8 mL PEN	three doses to be given in contres, Supplies, Nursing Services for brug only for facility administration. DOSE Inject 40 mg SC every week Inject 40 mg SC every other Inject 80 mg SC every other Inject 80 mg Day 1, followed week after initial dose Inject 40 mg SC every week Inject 40 mg SC every other Inject 40 mg SC every week Inject 40 mg SC every other Inject 80 mg SC every other	rolled setting. drug administration & DIRECTIONS week week by 40 mg every other week week week	n/therapy teach train.	QUANTITY/REFIL 28 days 84 days
Home Infusion/Contraction Prescriber's Office PRESCRIPTION Adalimumabaacf (Unbranded Idacio) Adalimumabadacio	nicade Biosimilars: First ram AIS: Diluents, Flushe e/Other Infusion Clinic: DON INFORMATION STRENGTH 40 mg/0.8 mL PEN	Inject 40 mg SC every week Inject 40 mg SC every week Inject 80 mg Day 1, followed week after initial dose Inject 40 mg SC every week Inject 80 mg SC every other Inject 80 mg SC every other Inject 80 mg SC every other week after initial dose Inject 40 mg SC every other Inject 80 mg SC every other	rolled setting. drug administration & DIRECTIONS week week by 40 mg every other week week week	n/therapy teach train.	QUANTITY/REFIL 28 days 84 days Refills:
Home Infusion/Contraction Prescriber's Office PRESCRIPTION Adalimumabaacf (Unbranded Idacio) Adalimumabadacio) Adalimumabadaz (unbranded Idacio)	nicade Biosimilars: First ram AIS: Diluents, Flushe e/Other Infusion Clinic: D ON INFORMATION STRENGTH 40 mg/0.8 mL PEN 40 mg/0.4 mL PEN 40 mg/0.4 mL PFS	Inject 40 mg SC every week Inject 40 mg SC every other Inject 40 mg SC every other Inject 80 mg Day 1, followed week after initial dose Inject 40 mg SC every week Inject 40 mg SC every other Inject 80 mg Day 1, followed week after initial dose Inject 40 mg SC every other Inject 80 mg SC on Day 1, follone week after initial dose	rolled setting. drug administration E & DIRECTIONS week week by 40 mg every other week week lowed by 40 mg ever	or week starting one	QUANTITY/REFIL 28 days 84 days Refills: Quantity:
Home Infusion/Control Prescriber's Office PRESCRIPTION Adalimumabaacf (Unbranded Idacio) Adalimumabadacio) Adalimumabadacio	nicade Biosimilars: First ram AIS: Diluents, Flushe e/Other Infusion Clinic: DON INFORMATION STRENGTH 40 mg/0.8 mL PEN	three doses to be given in contres, Supplies, Nursing Services for brug only for facility administration. DOSE Inject 40 mg SC every week Inject 40 mg SC every other Inject 80 mg SC every other Inject 80 mg Day 1, followed week after initial dose Inject 40 mg SC every other Inject 40 mg SC every other Inject 40 mg SC every other Inject 80 mg SC every other Inject 80 mg SC every other Inject 80 mg SC on Day 1, followe week after initial dose Inject 160 mg SC on Day 1 (si	rolled setting. drug administration E & DIRECTIONS week week by 40 mg every other week week lowed by 40 mg ever	h/therapy teach train. er week starting one ry other week starting er two consecutive	QUANTITY/REFIL 28 days 84 days Refills: Quantity: 28 days
Home Infusion/Control Prescriber's Office PRESCRIPTION Adalimumabaacf (Unbranded Idacio) Adalimumabadacio) Adalimumabadacio	nicade Biosimilars: First ram AIS: Diluents, Flushe e/Other Infusion Clinic: D ON INFORMATION STRENGTH 40 mg/0.8 mL PEN 40 mg/0.4 mL PEN 40 mg/0.4 mL PFS	Inject 40 mg SC every week Inject 40 mg SC every other Inject 40 mg SC every other Inject 80 mg SC every week Inject 40 mg SC every other Inject 80 mg SC every other Inject 80 mg SC every other Inject 40 mg SC every other Inject 80 mg SC every other Inject 80 mg SC every other Inject 40 mg SC every other Inject 40 mg SC every other Inject 80 mg SC every other Inject 80 mg SC on Day 1, follone week after initial dose Inject 160 mg SC on Day 1 (sidays), 80 mg on Day 15, then 40	rolled setting. drug administration E & DIRECTIONS week week by 40 mg every other week week lowed by 40 mg ever ngle-dose or split ov mg every week star	or week starting one ry other week starting er two consecutive ting on Day 29	QUANTITY/REFIL 28 days 84 days Refills: Quantity: 28 days 84 days
Home Infusion/Conferescriber's Office PRESCRIPTION Adalimumabaacf (Unbranded Idacio) Adalimumabaadz (unbranded unbranded Idacio)	nicade Biosimilars: First ram AIS: Diluents, Flushe e/Other Infusion Clinic: D ON INFORMATION STRENGTH 40 mg/0.8 mL PEN 40 mg/0.4 mL PEN 40 mg/0.4 mL PFS	Inject 40 mg SC every week Inject 40 mg SC every week Meek after initial dose Inject 80 mg SC every other Inject 40 mg SC every other Inject 40 mg SC every other Inject 80 mg SC every other Inject 80 mg SC every other Inject 40 mg SC every other Inject 80 mg SC every other Inject 80 mg SC on Day 1, followed week after initial dose Inject 160 mg SC on Day 1 (si days), 80 mg on Day 15, then 40 Inject 160 mg SC on Day 1 (si	rolled setting. drug administration E & DIRECTIONS week week by 40 mg every other week week lowed by 40 mg ever ongle-dose or split ov ong every week star ngle-dose or split ov	er week starting one er two consecutive ting on Day 29 er two consecutive	QUANTITY/REFIL 28 days 84 days Refills: Quantity: 28 days
Home Infusion/Control Prescriber's Office PRESCRIPTION Adalimumabaacf (Unbranded Idacio) Adalimumabadacio) Adalimumabadacio	nicade Biosimilars: First ram AIS: Diluents, Flushe e/Other Infusion Clinic: DON INFORMATION STRENGTH 40 mg/0.8 mL PEN 40 mg/0.4 mL PEN 40 mg/0.4 mL PFS (with needle guard)	Inject 40 mg SC every week Inject 40 mg SC every other Inject 40 mg SC every other Inject 80 mg SC every week Inject 40 mg SC every other Inject 80 mg SC every other Inject 80 mg SC every other Inject 40 mg SC every other Inject 80 mg SC every other Inject 80 mg SC every other Inject 40 mg SC every other Inject 40 mg SC every other Inject 80 mg SC every other Inject 80 mg SC on Day 1, follone week after initial dose Inject 160 mg SC on Day 1 (sidays), 80 mg on Day 15, then 40	rolled setting. drug administration E & DIRECTIONS week week by 40 mg every other week week lowed by 40 mg ever ongle-dose or split ov ong every week star ngle-dose or split ov	er week starting one er two consecutive ting on Day 29 er two consecutive	QUANTITY/REFIL 28 days 84 days Refills: Quantity: 28 days 84 days Refills:
Home Infusion/Coopersoriber's Office PRESCRIPTION Adalimumabased (Unbranded Idacio) Adalimumabased (Unbranded Idacio) Adalimumabased (Unbranded Idacio)	nicade Biosimilars: First ram AIS: Diluents, Flushe e/Other Infusion Clinic: D ON INFORMATION STRENGTH 40 mg/0.8 mL PEN 40 mg/0.4 mL PEN 40 mg/0.4 mL PFS	Inject 40 mg SC every week Inject 40 mg SC every week Meek after initial dose Inject 80 mg SC every other Inject 40 mg SC every other Inject 40 mg SC every other Inject 80 mg SC every other Inject 80 mg SC every other Inject 40 mg SC every other Inject 80 mg SC every other Inject 80 mg SC on Day 1, followed week after initial dose Inject 160 mg SC on Day 1 (si days), 80 mg on Day 15, then 40 Inject 160 mg SC on Day 1 (si	rolled setting. drug administration E & DIRECTIONS week week by 40 mg every other week week lowed by 40 mg ever ongle-dose or split ov ong every week star ngle-dose or split ov	er week starting one er two consecutive ting on Day 29 er two consecutive	QUANTITY/REFIL 28 days 84 days Refills: Quantity: 28 days 84 days Refills: Quantity: Quantity:
Home Infusion/Col Prescriber's Office PRESCRIPTION Adalimumabased (Unbranded dacio) Adalimumabased (Unbranded dacio) Adalimumabased (Unbranded dacio) Other	nicade Biosimilars: First ram AIS: Diluents, Flushe e/Other Infusion Clinic: DON INFORMATION STRENGTH 40 mg/0.8 mL PEN 40 mg/0.4 mL PEN 40 mg/0.4 mL PFS (with needle guard)	Inject 40 mg SC every week Inject 40 mg SC every week Meek after initial dose Inject 80 mg SC every week Inject 40 mg SC every other Inject 80 mg SC every other Inject 40 mg SC every other Inject 80 mg SC on Day 1, followed week after initial dose Inject 160 mg SC on Day 1 (si days), 80 mg on Day 15, then 40 mg SC on Day 1 (si days), 80 mg on Day 15, then 80 mg SC on D	rolled setting. drug administration A DIRECTIONS week week by 40 mg every other week lowed by 40 mg every mgle-dose or split ov mg every week star ngle-dose or split ov mg every other wee	er week starting one er two consecutive ting on Day 29 er two consecutive	QUANTITY/REFIL 28 days 84 days Refills: Quantity: 28 days 84 days Refills: Quantity: Refills:
Home Infusion/Col Prescriber's Office PRESCRIPTION MEDICATION Adalimumabased (Unbranded Idacio) Adalimumabased (Unbranded Idacio) Adalimumabased (Unbranded Idacio) Other Patient is interested in patient	nicade Biosimilars: First ram AIS: Diluents, Flushe e/Other Infusion Clinic: DON INFORMATION STRENGTH 40 mg/0.8 mL PEN 40 mg/0.4 mL PEN 40 mg/0.4 mL PFS (with needle guard) Strength:	Inject 40 mg SC every week Inject 40 mg SC every week Meek after initial dose Inject 80 mg SC every other Inject 80 mg SC every other Inject 40 mg SC every other Inject 80 mg SC every other Inject 40 mg SC every other Inject 40 mg SC every other Inject 80 mg SC every other Inject 80 mg SC every other Inject 80 mg SC on Day 1, followed week after initial dose Inject 160 mg SC on Day 1 (sidays), 80 mg on Day 15, then 40 mg SC on Day 1 (sidays), 80 mg on Day 15, then 80 mg SC on Day 1 (sidays), 80 mg on Day 15, then 80 mg SC on Day 1.	rolled setting. drug administration E & DIRECTIONS week week by 40 mg every other week lowed by 40 mg ever ngle-dose or split ov mg every week star ngle-dose or split ov mg every week star ngle-dose or split ov mg every week star	er week starting one er two consecutive ting on Day 29 er two consecutive ek starting on Day 29	QUANTITY/REFIL 28 days 84 days Refills: Quantity: 28 days 84 days Refills: Quantity: Refills:
Home Infusion/Control Prescriber's Office PRESCRIPTION Adalimumabaacf (Unbranded Idacio) Adalimumabadaz (unbranded Version of Hyrimoz) Other Patient is interested in patient PRESCRIBER \$	nicade Biosimilars: First ram AIS: Diluents, Flushe e/Other Infusion Clinic: DON INFORMATION STRENGTH 40 mg/0.8 mL PEN 40 mg/0.4 mL PEN 40 mg/0.4 mL PFS (with needle guard) Strength: t support programs SIGNATURE REQUIRI	three doses to be given in contrest. Supplies, Nursing Services for brug only for facility administration. DOSE Inject 40 mg SC every week Inject 40 mg SC every other Inject 80 mg SC every other Inject 80 mg SC every week Inject 40 mg SC every other Inject 80 mg SC on Day 1, followed week after initial dose Inject 80 mg SC on Day 1, followed week after initial dose Inject 160 mg SC on Day 1 (sidays), 80 mg on Day 15, then 40 mg SC, so Day 1 (sidays), 80 mg on Day 15, then 80 mg SC on Day 1 (sidays), 80 mg on Day 15, then 80 mg SC on Day 1 (sidays), 80 mg on Day 15, then 80 mg SC on Day 1 (sidays), 80 mg on Day 15, then 80 mg SC on Day 1 (sidays), 80 mg S	rolled setting. drug administration E & DIRECTIONS week week by 40 mg every other week lowed by 40 mg ever ongle-dose or split ov ong every week star ngle-dose or split ov ong every other wee Ancill OT ALLOWED)	er week starting one er two consecutive ting on Day 29 er two consecutive ek starting on Day 29 lary supplies and kits provided as need	QUANTITY/REFIL 28 days 84 days Refills: Quantity: 28 days 84 days Refills: Quantity: Refills:
Home Infusion/Co *Prescriber's Office PRESCRIPTION Adalimumab- aacf (Unbranded Idacio) Adalimumab- adaz (unbranded version of Hyrimoz) Patient is interested in patient PRESCRIBER \$	nicade Biosimilars: First ram AIS: Diluents, Flusher e/Other Infusion Clinic: DON INFORMATION STRENGTH 40 mg/0.8 mL PEN 40 mg/0.4 mL PEN 40 mg/0.4 mL PFS (with needle guard) Strength: tsupport programs BIGNATURE REQUIRE / Brand Medically Necessary / Do	Inject 40 mg SC every week Inject 40 mg SC every other Inject 40 mg SC every other Inject 40 mg SC every other Inject 80 mg Day 1, followed week after initial dose Inject 80 mg SC every other Inject 80 mg SC every other Inject 80 mg SC every other Inject 40 mg SC every other Inject 40 mg SC every other Inject 80 mg SC on Day 1, followed week after initial dose Inject 160 mg SC on Day 1 (si days), 80 mg on Day 15, then 40 mg SC on Day 1 (si days), 80 mg on Day 15, then 80 mg SC on Day 1 (si days), 80 mg on Day 15, then 80 mg Dose:	rolled setting. drug administration E & DIRECTIONS week week by 40 mg every other week lowed by 40 mg ever ngle-dose or split ov mg every week star ngle-dose or split ov mg every week star ngle-dose or split ov mg every week star	er week starting one er two consecutive ting on Day 29 er two consecutive ek starting on Day 29 lary supplies and kits provided as need	QUANTITY/REFIL 28 days 84 days Refills: Quantity: 28 days 84 days Refills: Quantity: Refills:
Home Infusion/Co *Prescriber's Office PRESCRIPTION Adalimumabaacf (Unbranded Idacio) Adalimumabadaz (unbranded version of Hyrimoz) Other PRESCRIBER S "Dispense As Written" DAW / May Not Substit	nicade Biosimilars: First ram AIS: Diluents, Flusher e/Other Infusion Clinic: DON INFORMATION STRENGTH 40 mg/0.8 mL PEN 40 mg/0.4 mL PEN 40 mg/0.4 mL PFS (with needle guard) Strength: tsupport programs BIGNATURE REQUIRE / Brand Medically Necessary / Do	Inject 40 mg SC every week Inject 40 mg SC every other Inject 80 mg SC every week Inject 40 mg SC every other Inject 80 mg SC every other Inject 40 mg SC every other Inject 40 mg SC every other Inject 40 mg SC every other Inject 80 mg SC every other Inject 80 mg SC on Day 1, followed week after initial dose Inject 80 mg SC on Day 1, followed week after initial dose Inject 160 mg SC on Day 1 (sidays), 80 mg on Day 15, then 80 mg SC on Day 1 (sidays), 80 mg on Day 15, then 80 mg SC on Day 1 (sidays), 80 mg on Day 15, then 80 mg SC on Day 1 (sidays), 80 mg on Day 15, then 80 mg SC on Day 1 (sidays), 80 mg SC on Da	rolled setting. drug administration n E & DIRECTIONS week week by 40 mg every othe week lowed by 40 mg ever ngle-dose or split ov 0 mg every week star ngle-dose or split ov 0 mg every other wee DT ALLOWED) May Substitute / Product	ry other week starting one er two consecutive ting on Day 29 er two consecutive ek starting on Day 29 lary supplies and kits provided as need	QUANTITY/REFIL 28 days 84 days Refills: Quantity: 28 days 84 days Refills: Quantity: Refills:

		Patient , Prescriber an			
			Patient		
Prescriber Name: _ Patient Clinical Info		Prescribe	er Phone:		
Allergies:					
Weight:		In/cm TB T	est Result:	Date:	
	ON INFORMATION				_
MEDICATION	STRENGTH	DOSE &	directions	QUANTITY/REF	FILLS
Adalimumab- fkjp (unbranded version of Hulio)	☐ 40 mg/0.8 mL PFS ☐ 40 mg/0.8 mL PEN	☐ Inject 40 mg SC every week☐ Inject 40 mg SC every other v☐ Inject 80 mg SC every other v☐ Inject 80 mg SC on Day 1, folloone week after initial dose☐ Inject 160 mg SC on Day 1 (sir days), 80 mg on Day 15, then 40☐ Inject 160 mg SC on Day 1 (sir days), 80 mg on Day 15, then 80	veek veek owed by 40 mg every other we ogle-dose or split over two cons mg every week starting on Day ogle-dose or split over two cons	Quantity: 28 days secutive 84 days 729 Refills:	
Amjevita (adalimumab-atto)	☐ 40 mg/0.8 mL PFS ☐ 40 mg/0.8 mL PEN	☐ Inject 40 mg SC every week☐ Inject 40 mg SC every other v☐ Inject 80 mg SC every other v☐ Inject 80 mg Day 1, followed bweek after initial dose☐ Inject 160 mg SC on Day 1 (given consecutive days), 80 mg on Day other week dosing two weeks lat	veek veek by 40 mg every other week star ven in one day or split over two v 15. Begin 40 mg weekly or 80	Quantity: Tting one 28 days 84 days Refills:	
Avsola	100 mg vial	☐ Induction Dose: Infuse IV at 5 0, 2, 6 and every 8 weeks therea: ☐ Maintenance Dose: Infuse IV (Dose =mg) every 8 weeks	mg/kg (Dose =mg) at we ter	eks Quantity: # of 100 mg vial(s) Refills:)
☐ Bimzelx	☐ 2 x 160 mg/mL PEN☐ 2 x 160 mg/mL PFS	Loading Dose: ☐ Inject 320 mg (2 x 160 mg/ml Maintenance Dose: ☐ Inject 320 mg (2 x 160 mg/ml Patients ≥ 120 kg (264lbs) may co ☐ Inject 320 mg (2 x 160 mg/ml	.) SC every 8 weeks onsider:	Loading Dose: Quantity: 28 Days Refills: 4 Maintenance Dose Quantity: 28 Days 28 Days 56 Days Refills:	<u>e</u> :
☐ Cimzia	Cimzia Starter Kit (6 prefilled syringes)	Psoriasis Loading Dose: 400 mg (given as 2 subcutan other week Patients (with body weight ≤ 2 subcutaneous injections of 200 2 and 4, followed by 200 mg eve Psoriatic Arthritis Loading Dose: 400 mg (given as 2 subcutan and at week 2 and 4, followed by	90 kg): 400 mg (given as 9 mg each) initially and at week ry other week eous injections of 200 mg each	Quantity: 1 Kit Refills: 0	
☐ Cimzia	200 mg/1 mL prefilled syringe	Psoriasis Maintenance Dose: 400 mg (given as 2 subcutaneous injections of 200 mg each) every other week 200 mg every other week Psoriatic Arthritis Maintenance Dose: 200 mg every other week 400 mg (given as 2 subcutaneous injections of 200 mg each) every 4 weeks		Quantity: Refills:	
Other	Strength:	Dose:		Quantity: Refills:	
Patient is interested in pa		STAMP SIGNATURE NOT ALLOWED		es and kits provided as needed for administ	stration
PRESCRIBER 1	SIGNATURE REQU	IRED (STAMP SIGNATU	RE NOT ALLOWED)		
/ May Not Substitute Prescriber's Signa	ature:	Not Substitute / No Substitution / DAW Date: pr writes the words "No Substitution"	May Substitute / Product Selection Substitution Permissible Prescriber's Signature:		

	Please Complete Patic	ent , Prescriber an	d Patient Clini	cal Information	
Patient Name:	Pa	atient DOB:		Patient Phone:	
		Prescribe	er Phone:		
Patient Clinical Informa					
Allergies:	lb/kg Height:	In /one TD T	ant Donults	Datas	
5 PRESCRIPTION	tb/kg Height:	IN/CM 1B I	est Result:	Date:	
			OOE O DIDECTION	10	OLIANTITY/DEELLO
MEDICATION	STRENGTH		OSE & DIRECTION	15	QUANTITY/REFILLS
☐ Cosentyx	☐ 75 mg/0.5 mL PFS ☐ 150 mg/mL PEN ☐ 150 mg/mL PFS ☐ 150 mg/mL PEN ☐ 150 mg/mL PFS ☐ 300 mg/2 mL PEN	Loading Dose: Inject 75 mg SC on Inject 150 mg SC or Inject 300 mg SC or Maintenance Dose: Inject 75 mg SC or Inject 75 mg SC or Inject 150 mg SC or Inject 150 mg SC or Inject 300 mg SC or Inject 300 mg SC or Inject 300 mg SC or	Neeks 0, 1, 2, 3 Neeks 0, 1, 2, 3 Week 4, then every 4 ery 4 weeks Neek 4, then every ery 4 weeks Neek 4, then every very 4 weeks	4 weeks thereafter	Loading Dose: Quantity: 28 days Refills: 0 Maintenance Dose: Quantity: 28 days Refills:
☐ Dupixent	☐ PFS 300 mg/2 mL prefilled syringe ☐ Pen* 300 mg/2 mL prefilled pen *Comes in cartons of 2	Initial Prurigo Nodularis Inject 600 mg SC (2 SC every other week Maintenance Prurigo N Inject 300 mg SC every	s <u>Dose:</u> 2-300 mg injections) Iodularis Dose:	initially then 300 mg	Quantity: 28-day supply 84-day supply Other:Day supply Refills: 1 year Other:Refills
☐ Enbrel	50 mg/mL Mini 50 mg/mL PEN 50 mg/mL PFS 25 mg/0.5 mL PFS 25 mg/0.5 mL Vial	Loading Dose: Inject 50 mg SC twi 3 months, then mainter Maintenance Dose: Inject 50 mg SC one Inject mg Sc	nance dosing ce weekly	ys apart) for	Loading Dose: Quantity: 84 days Refills: 0 Maintenance Dose: Quantity: 28 days Refills:
☐ Hadlima	☐ 40 mg/0.4 mL PEN☐ 40 mg/0.8 mL PEN☐ 40 mg/0.4 mL PFS☐ 40 mg/0.8 mL PFS	☐ Inject 40 mg SC eve ☐ Inject 40 mg SC eve ☐ Inject 80 mg SC eve ☐ Inject 80 mg SC on week starting one weel ☐ Inject 160 mg SC or consecutive days), 80 starting on Day 29 ☐ Inject 160 mg SC or consecutive days), 80 week starting on Day 2	ery other week ery other week Day 1, followed by 40 k after initial dose n Day 1 (single-dose omg on Day 15, then 4 n Day 1 (single-dose omg on Day 15, then 8	or split over two 00 mg every week or split over two	Quantity: 28 days 84 days Refills:
Other	Strength:	Dose:			Quantity: Refills:
Patient is interested in patient: PRESCRIBER SIG	support programs STAMPS GNATURE REQUIRED (SIGNATURE NOT ALLOWED			ded as needed for administration
"Dispense As Written" / Brar / May Not Substitute Prescriber's Signatur	nd Medically Necessary / Do Not Substi	itute / No Substitution / DAW	May Substitute / Produ Substitution Permissib Prescriber's Sign	uct Selection Permitted /	Date:submit electronic prescription

	Please Comple	ete Patient , Prescriber an	d Patient Clinical Informatio	<u>n</u>
		Patient DOB:	Patient Phone:	
	ne:	Prescribe	r Phone:	
Patient Clinical				
Allergies:	lb/kg Height	In /are TD T		
			est Result: Date: _	
	TION INFORMATION			
MEDICATION	STRENGTH		DIRECTIONS	QUANTITY/REFILLS
☐ Hulio	☐ 40 mg/0.8 mL PFS ☐ 40 mg/0.8 mL PEN	one week after initial dose Inject 160 mg SC on Day 1 (single days), 80 mg on Day 15, then 40 mg Inject 160 mg SC on Day 1 (single days), 80 mg on Day 15, then 80 mg	ek ed by 40 mg every other week starting e-dose or split over two consecutive	Quantity: 28 days 84 days Refills:
☐ Humira	☐ 40 mg/0.4 mL PFS ☐ 40 mg/0.4 mL Pen ☐ 80 mg/0.8 mL PFS ☐ 80 mg/0.8 mL Pen	subsequent doses Inject 160 mg SC on Day 1 (single days), 80 mg on Day 15, then 40 mg Inject 160 mg SC on Day 1 (single	ek O mg every other week on day 8 and e-dose or split over two consecutive	28 days 84 days Refills:
☐ Hyrimoz	40 mg/0.4 mL PEN 80 mg/0.8 mL PEN 40 mg/0.4 mL PFS (with needle guard) Psoriasis Starter Kit (1-80 mg and 2-40 mg PEN)	one week after initial dose Inject 160 mg SC on Day 1 (single days), 80 mg on Day 15, then 40 mg Inject 160 mg SC on Day 1 (single	ek ed by 40 mg every other week starting e-dose or split over two consecutive	28 days 84 days Refills:
☐ Idacio	☐ 40 mg/0.8 mL PEN ☐ 40 mg/0.8 mL PFS	☐ Inject 40 mg SC every week☐ Inject 40 mg SC every other wee☐ Inject 80 mg SC every other wee☐ Inject 80 mg SC on Day 1, follow one week after initial dose		28 days 84 days Refills:
□ Ilumya	100 mg/mL prefilled syringe	Psoriasis Induction Dose: Inject one pre-filled syringe (100 mg) SC at weeks 0 and 4, then maintenance dosing. Psoriasis Maintenance Dose: Inject one pre-filled syringe (100 mg) SC every 12 weeks.		Quantity: Refills:
☐ Inflectra		☐ Induction Dose: Infuse IV at 5 mg	g/kg	Quantity:
☐ Infliximab	100 mg vial	(Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 8 weeks		# of 100 mg vial(s) Refills:
Other	Strength:	Dose:		Quantity: Refills:
	d in patient support programs BER SIGNATURE REQ	STAMP SIGNATURE NOT ALLOWED UIRED (STAMP SIGNATURE)	, , , ,	ovided as needed for administration
/ May Not Substitu Prescriber's \$	te Signature:	Do Not Substitute / No Substitution / DAW Date: riber writes the words "No Substitution"	May Substitute / Product Selection Permitted / Substitution Permissible Prescriber's Signature:ATTN: New York and Iowa providers, pleas	

	Please Compl	ete Patient , Prescriber an	d Patient Clinical Information	
Patient Name: _			Patient Phone:	
		Prescribe	r Phone:	
Patient Clinical				
Allergies:	lb/kg Heigh	nt: In/cm TBTe	est Result: Date:	
	ION INFORMATION			
MEDICATION	STRENGTH	DOSE	& DIRECTIONS	QUANTITY/REFILL
Litfulo	50 mg capsule	☐ Take 50 mg orally once daily with	or without food	28 days 84 days Refills:
Olumiant	2 mg tablet 4 mg tablet	2 mg PO once daily 4 mg PO once daily		Quantity: Refills:
Orencia	125 mg/mL prefilled syringe	Inject 125 mg SC once weekly		Quantity: Refills:
☐ Otezla	Titration Starter Pack	Day 1: 10 mg PO in the morning. Day 2: 10 mg PO in the morning and 10 mg PO in the evening. Day 3: 10 mg PO in the morning and 20 mg PO in the evening. Day 4: 20 mg PO in the morning and 20 mg PO in the evening. Day 5: 20 mg PO in the morning and 30 mg PO in the evening. Day 6 and thereafter: 30 mg PO twice daily.		Quantity: 1 pack Refills: 0
Otezla	30 mg tablet	Maintenance Dose: 30 mg tablet	PO twice daily.	Quantity: Refills:
Remicade Renflexis	100 mg vial	☐ Induction Dose: Infuse IV at 5 mg/ (Dose =mg) at weeks 0, 2, 6 and ☐ Maintenance Dose: Infuse IV at 5		Quantity: # of 100 mg vial(s) Refills:
Rinvoq	15 mg	Take one 15 mg tablet PO daily		Quantity: Refills:
Siliq	Carton of two 210 mg/1.5 mL single-dose prefilled syringes	Inject one prefilled syringe (210 mg) SC at weeks 0, 1 and 2, followed by one prefilled syringe (210 mg) every 2 weeks. Prescribers must be certified in the SILIQ REMS Program to prescribe SILIQ. Please visit the following REMS website to register before prescribing SILIQ: SILIQ REMS Website (https://siliqrems.com/SiliqUI/home.u)		Quantity: Refills:
Simponi	50 mg/0.5 mL SmartJect Autoinjector 50 mg/0.5 mL prefilled syringe	Psoriatic Arthritis Dose: Inject 50 mg SC once a month.		Quantity: Refills:
☐ Simponi ARIA	50 mg/4 mL in a single- dose vial	Psoriatic Arthritis Dosing: Induction Dose: 2 mg/kg IV infusion over 30 minutes at weeks 0 and 4, then every 8 weeks thereafter Maintenance Dose: 2 mg/kg IV infusion over 30 minutes every 8 weeks		Quantity: # of 50 mg vial Refills:
Skyrizi	☐ 150 mg/mL single-dose Pen ☐ 150 mg/mL single-dose prefilled syringe	Psoriasis Induction Dose: Inject 150 mg SC at Weeks 0 and 4, then maintenance dosing. Psoriasis Maintenance Dose: Inject 150 mg SC every 12 weeks.		Quantity: Refills:
Sotyktu	6 mg tablet	Take one 6 mg tablet PO once daily		Quantity: Refills:
Other	Strength:	□ Dose:		Quantity: Refills:
	in patient support programs ER SIGNATURE REC	STAMP SIGNATURE NOT ALLOWED QUIRED (STAMP SIGNATUR	Ancillary supplies and kits provided RE NOT ALLOWED)	as needed for administration
/ May Not Substitut Prescriber's S	ignature:	/ Do Not Substitute / No Substitution / DAW	May Substitute / Product Selection Permitted / Substitution Permissible Prescriber's Signature:	Date:
CA, MA, NC & PR:	Interchange is mandated unless Pres	scriber writes the words "No Substitution"	ATTN: New York and Iowa providers, please subm	nit electronic prescription

	Please Complete	e Patient , Prescriber a	and Patient Clini	cal Information	<u>h</u>
		Patient DOB:		Patient Phone: _	
		Prescr	iber Phone:		
Patient Clinical I					
Allergies:	lb/kg Height: _	In/cm TE	P Toet Pocult:	Date:	
DDESCDID	TION INFORMATION	III/OIII 11	3 1651 Result	Date	
MEDICATION	STRENGTH	DOS	E & DIRECTIONS		QUANTITY/REFILLS
MEDICATION	JIKENGIII	PsO Peds patients (6 to 17yo):			QUARITI IZ RELIES
☐ Stelara	☐ 45 mg/0.5 mL prefilled syringe ☐ 90 mg/mL prefilled syringe	< 60 kg: Inject 0.75 mg/kg: thereafter. 60 kg to 100 kg: Inject 45 m 12 weeks thereafter. > 100 kg: Inject 90 mg SC at thereafter. > 100 kg: Inject 90 mg SC at thereafter. > 60 kg: Inject 0.75 mg/kg 12 weeks thereafter. > 60 kg: Inject 45 mg SC at thereafter. > 100 kg with co-existent n weeks 0 and 4, then every 12 vests a later, followed by 45 mesh 12 weeks later, followed by 90 mesh 2 mesh 100 kg (220lbs) with co-esc weeks 0 and 4, then every 100 kg (220lbs) with co-esc weeks 0 and 4, then every 100 kg (220lbs) with co-esc weeks 0 and 4, then every 100 kg (220lbs) with co-esc weeks 0 and 4, then every 100 kg (220lbs) with co-esc weeks 0 and 4, then every 100 kg (220lbs) with co-esc weeks 0 and 4, then every 100 kg (220lbs) with co-esc weeks 0 and 4, then every 100 kg (220lbs) with co-esc weeks 0 and 4, then every 120 kg (220lbs) with co-esc weeks 0 and 4, then every 100 kg (220lbs) with co-esc weeks 0 and 4, then every 100 kg (220lbs) with co-esc weeks 0 and 4, then every 100 kg (220lbs) with co-esc weeks 0 and 4, then every 100 kg (220lbs) with co-esc weeks 0 and 4, then every 100 kg (220lbs) with co-esc weeks 0 and 4, then every 100 kg (220lbs) with co-esc weeks 0 and 4, then every 100 kg (220lbs) with co-esc weeks 0 and 4, then every 100 kg (220lbs) with co-esc weeks 0 and 4, then every 100 kg (220lbs) with co-esc weeks 0 kg (220lbs	at weeks 0 and 4, then early seeks thereafter. Okg (220 lbs): Inject 45 mg every 12 weeks. Okg (220 lbs): Inject 90 mg every 12 weeks. Oand 4, then every 12 weeks. Oand 4, then every 12 weeks.	every 12 weeks hen every very 12 weeks 90 mg SC at mg SC initially and mg SC initially and	Quantity: Refills:
☐ Taltz	☐ 80 mg Single Dose Autoinjector ☐ 80 mg Single Dose prefilled syringe	Psoriasis Dosing: Starting Dose: Inject SC tw. first induction dose 2 weeks la Induction Dose: Inject SC c 2-10). Final Induction Dose: Inject SC c Maintenance Dose: Inject SC pediatric Psoriasis Dosing: For patients weighing less that 40 mg at Week 0, followed For patients weighing 25-50 k 80 mg at Week 0, followed For patients weighing greater 160 mg (two 80 mg injectic weeks	to 80 mg injections on Dater. The second of	ry 2 weeks (weeks n (week 12). every 4 weeks. ks.	Quantity: 3 Pens/Syringes 2 Pens/Syringes 1 Pen/Syringe Refills:
☐ Taltz	80 mg Single Dose Autoinjector 80 mg Single Dose prefilled syringe	Psoriatic Arthritis Dosing: Starting Dose: Inject SC tw. Maintenance Dose: Inject S			Quantity: Refills:
Other	Strength:	Dose:			Quantity: Refills:
	in patient support programs ER SIGNATURE REQU	STAMP SIGNATURE NOT ALLOWED IRED (STAMP SIGNAT		, ,,	vided as needed for administration
DAW / May Not Sub Prescriber's Si	en" / Brand Medically Necessary / Do ostitute ignature: interchange is mandated unless Prescribe	Date:	May Substitute / Product Substitution Permissible Prescriber's Signat ATTN: New York a	ture:	Date:

	Please Comple	te Patient , Prescriber a	<u>nd Patient Clinical Informatio</u>	on .
Patient Name: _		Patient DOB:	Patient Phone:	
Prescriber Nam	me: Prescriber Phone:			
<u>Patient Clinical I</u>				
Allergies:			Test Result: Date:	
			Test Result: Date:	
	TION INFORMATION			
MEDICATION	STRENGTH	DOSE	& DIRECTIONS	QUANTITY/REFILLS
☐ Tremfya	☐ 100 mg/mL prefilled syringe ☐ 100 mg/mL One-Press patient-controlled injector	Starting Dose: Inject 100 mg Starting Maintenance Dose: Inject 100	C at weeks 0 and 4, then maintenance mg SC every 8 weeks	Quantity: Refills:
Xeljanz	☐ 5 mg tablet☐ 11 mg XR tablet	☐ Take one 5 mg tablet PO twice☐ Take one 11 mg PO once daily	e daily	Quantity: Refills:
☐ Yuflyma	40 mg/0.4 mL PEN 40 mg/0.4 mL PFS 40 mg/0.4 mL PFS (with safety guard) 80 mg/0.8 mL PEN	one week after initial dose Inject 160 mg SC on Day 1 (sind days), 80 mg on Day 15, then 40 mInject 160 mg SC on Day 1 (sind loss)	eek wed by 40 mg every other week starting gle-dose or split over two consecutive	28 days 84 days Refills:
Other	Strength:	Dose:		Quantity: Refills:
	in patient support programs	STAMP SIGNATURE NOT ALLOWED		rovided as needed for administration
		UIRED (STAMP SIGNATU		
DAW / May Not Sub	ostitute	Do Not Substitute / No Substitution / Date:	May Substitute / Product Selection Permitted / Substitution Permissible Prescriber's Signature:	Date:

Dermatology Enrollment Form Nursing Orders

Ple	ease Comp	olete Patient , Prescriber and Patient Clinical Informati	on
Patient Name:		Patient DOB:Patient Phone	•
Prescriber Name:		Prescriber Phone:	
<u>Patient Clinical Informatio</u>	<u>on:</u>		
Allergies:			
		ht: In/cm TB Test Result: Date	
		ON **ITEMS BELOW THIS LINE WILL ONLY BE SENT FOR INFUSIONS DO	
MEDICATION/SUPPLIES	ROUTE	DOSE /STRENGTH/ DIRECTIONS	QUANTITY/REFILLS
Catheter: PIV PORT CVC/PICC	IV	Catheter Care/Flush – Only on drug admin days – SASH or PRN to maintain IV access and patency PIV: NS 5 mL (Heparin 10 units/mL 3-5 mL if multiple days) CVC/PICC: NS 10 mL & Heparin 10 units/mL or 100 units/mL 3-5 mL PORT: 10 mL sterile saline to access PORT w/ huber needle NS 10 mL & Heparin 100 units/mL 3-5 mL	Quantity: Refills:
Hydration:	IV	Pre:	Hydration max infusion rate mL/hr (Adult max rate 250 mL/hr unless otherwise indicated)
☐Epinephrine **nursing requires**	□ IM □ SC	☐ 1:1000, 0.3 mg/0.3 mL (greater than 30 kg/66lbs) ☐ 1:1000, 0.15 mg/0.3 mL (15-30 kg/33-66lbs) ☐ 1:1000, 0.01 mg/kg, Max 0.3 mg (under 15 kg) Mild-Moderate Reactions. May repeat in 3-5 minutes as needed For severe allergic reaction also call 911	Quantity: Refills:
Diphenhydramine Oral	РО	Premedication: ☐ 12.5 mg/kg (0-30 kg) ☐ 25 mg ☐ 50 mg (Over 30 kg)	Quantity: Refills:
Diphenhydramine 50 mg/mL vial **nursing required**	Slow IV	1 mg/kg (under 15 kg) 12.5 mg-50 mg (15-30 kg) 25 mg-50 mg (Over 30 kg) If mild/moderate reaction: may repeat in 3-5 minutes as needed (Adult max dose: 100 mg/day) If severe allergic reaction: call 911	Quantity: Refills:
☐ Flush Orders:	Peripheral Access Central Venous Access	☐ 10 mL NS post flush ☐ 50 mL NS post flush (recommended if no post-hydration) ☐ Other:	Send quantity sufficient for medication days supply
Additional Medication:			
Patient is interested in patient supp PRESCRIBER SIGN		STAMP SIGNATURE NOT ALLOWED Ancillary supplies and kits EQUIRED (STAMP SIGNATURE NOT ALLOWED)	Drovided as needed for administration
DAW / May Not Substitute Prescriber's Signature:		y / Do Not Substitute / No Substitution / Date: Date: Base of the words "No Substitution" May Substitute / Product Selection Permitted / Substitution Permissible Prescriber's Signature: ATTN: New York and Iowa providers, place of the words "No Substitution" ATTN: New York and Iowa providers, place of the words "No Substitution"	Date:

The information provided above is true and accurate to the best of my knowledge, with supporting documentation in the patient's medical record. By signing above, I hereby authorize CVS Specialty Pharmacy and/or its affiliate pharmacies to complete and submit prior authorization (PA) requests to payors for the prescribed medication for this patient and to attach this Enrollment Form to the PA request as my signature.

CONFIDENTIALITY NOTICE: This communication and any attachments may contain confidential and/or privileged information for the use of the designated recipients named above. If you are not the intended recipient, you are hereby notified that you have received this communication in error and that any review, disclosure, dissemination, distribution or copying of it or its contents is prohibited. If you have received this communication in error, please notify the sender immediately by telephone and destroy all copies of this communication and any attachments.

Plan member privacy is important to us. Our employees are trained regarding the appropriate way to handle members' private health information.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Specialty and/or one of its affiliates